Author index

Abd El Ghafar, N.Y. *see* Aly, M.M.
Ádám, A.L., Galal, A.A., Manninger, K. & Barna, B. Inhibition of the development of leaf rust (*Puccinia recondita*) by treatment of wheat with allopurinol and production of a hypersensitive-like reaction in a compatible host, 317
Adams, A.N. *see* Davies, D.L.
Adams, M.J. *see* Chen, J.
Adenle, V.O. & Cardwell, K.F. Seed transmission of maize downy mildew (*Peronosclerospora sorghi*) in Nigeria, 628
Adorada, D.L., Biles, C.L., Liddell, C.M., Fernández-Pavía, S., Waugh, K.O. & Waugh, M.E. Disease development and enhanced susceptibility of wounded pepper roots to *Phytophthora capsici*, 719
Aguilar, I., Sánchez, F. & Ponz, F. Different forms of interference between two tobamoviruses in two different hosts, 659
Aharoni, Y. *see* Fallik, E.
Akimoto, T. *see* Murakami, H.
Alfonso, C., Raposo, R. & Melgarejo, P. Genetic diversity in *Botrytis cinerea* populations on vegetable crops in greenhouses in south-eastern Spain, 243
Al-Hassani, H.A. *see* El-Tarabily, K.A.
Aly, M.M. & Abd El Ghafar, N.Y. Bacterial wilt of artichoke caused by *Ralstonia solanacearum* in Egypt, 806
Andrés, M.F., Pinochet, J., Hernández-Dorrego, A. & Delibes, A. Detection and analysis of inter- and intraspecific diversity of *Pratylenchus* spp. using isozyme markers, 640
Andrivon, D. *see* Bouchek-Mechiche, K.
Andrivon, D. *see* Hélias, V.
Andrivon, D. *see* Hélias, V.
Annamalai, P. *see* Hsu, Y.H.
Aramburu, J.M. & Rovira, M. Incidence and natural spread of apple mosaic ilarvirus in hazel in north-east Spain, 423
Asher, M.J.C. *see* Mutasa-Gottgens, E.S.
Asher, M.J.C. *see* Webb, C.R.
Ayllón, M.A. *see* d'Urso, F.

Ballester-Olmos, J.F. *see* Galipienso, L.
Barna, B. *see* Ádám, A.L.
Barrault, G. *see* Roustaee, A.
Barrault, G. *see* Roustaee, A.
Bastidas, O. *see* Vega-Sánchez, M.E.
Bateman, G.L. *see* Dawson, W.A.J.M.
Béasse, C., Ney, B. & Tivoli, B. A simple model of pea (*Pisum sativum*) growth affected by *Mycosphaerella pinodes*, 187
Bélanger, R.R. *see* Ongena, M.
Benhamou, N. *see* Ongena, M.
Bertrand, B., Nuñez, C. & Sarah, J.-L. Disease complex in coffee involving *Meloidogyne arabicida* and *Fusarium oxysporum*, 383
Biles, C.L. *see* Adorada, D.L.
Blanche, K.R. *see* Tran-Nguyen, L.
Blaschke, H. *see* Jung, T.
Børja, I. *see* Moricca, S.
Bouchek-Mechiche, K., Pasco, C., Andrivon, D. & Jouan, B. Differences in host range, pathogenicity to potato cultivars and response to soil temperature among *Streptomyces* species causing common and netted scab in France, 3
Brunt, A.A. & Spence, N.J. The natural occurrence of *hibiscus chlorotic ringspot virus* (*Carmovirus; Tombusviridae*) in aibika or bele (*Abelmoschus manihot*) in some South Pacific Island countries, 798
Brurberg, M.B. *see* Hermansen, A.
Burdon, J.J. *see* Thrall, P.H.
Burdon, J.J. *see* Wellings, C.R.
Burgess, T. *see* Jackson, T.J.
Burritt, D.J. *see* Clarke, S.F.
Butterworth, P.J. *see* Harvey, P.R.

Capretti, P. *see* Santini, A.
Cardwell, K.F. *see* Adenle, V.O.
Carter, J.P., Rezanoor, H.N., Desjardins, A.E. & Nicholson, P. Variation in *Fusarium graminearum* isolates from Nepal associated with their host of origin, 452
Caten, C.E. & Newton, A.C. Variation in cultural characteristics, pathogenicity, vegetative compatibility and electrophoretic karyotype within field populations of *Stagonospora nodorum*, 219
Chang, K.F. *see* Wang, H.
Chang, W.C. *see* Hsu, Y.H.
Chen, J., Chen, J.-P., Yang, J.-P., Cheng, Y., Diao, A., Adams, M.J. & Du, J. Differences in cultivar response and complete sequence analysis of two isolates of wheat yellow mosaic bymovirus in China, 370
Chen, J.-P. *see* Chen, J.
Chen, Y.Y. *see* Hsu, Y.H.
Cheng, Y. *see* Chen, J.
Chillet, M. *see* De Lapeyre De Bellaire, L.
Chwarszczynska, D.M. *see* Mutasa-Gottgens, E.S.
Ciuffo, M. *see* Roggero, P.
Clarke, S.F., Burritt, D.J., Jameson, P.E. & Guy, P.L. Effects of plant hormones on white clover mosaic potexvirus double-stranded RNA, 428
Clarkson, J.P., Kennedy, R. & Phelps, K. The effect of temperature and water potential on the production of conidia by sclerotia of *Botrytis squamosa*, 119
Cloquemin, G. *see* Seddas, A.
Collin, H.A. *see* Dugdale, L.J.
Collinge, D.B. *see* Madriz-Ordeñana, K.
Colquhoun, I. *see* Jackson, T.J.
Colson-Hanks, E.S. & Deverall, B.J. Effect of 2,6-dichloroisonicotinic acid, its formulation materials and benzothiadiazole on systemic resistance to alternaria leaf spot in cotton, 171
Colvin, J. *see* Zhang, X.-S.
Cook, S. *see* Hsiang, T.
Copel, A. *see* Fallik, E.
Costes, S. *see* Roustaee, A.
Cowger, C., Hoffer, M.E. & Mundt, C.C. Specific adaptation by *Mycosphaerella graminicola* to a resistant wheat cultivar, 445
Cronjé, P., Dabek, A.J., Jones, P. & Tymon, A.M. First report of a phytoplasma associated with a disease of date palms in North Africa, 801
Cronjé, P., Dabek, A.J., Jones, P. & Tymon, A.M. Slow decline: a new disease of mature date palms in North Africa associated with a phytoplasma, 804
Curir, P., Danieli, B., Dolci, M., Pasini, C., Guglieri, L. & Sacco, M. Reductive detoxication of the acetophenone skeleton of the carnation phytoanticipin by *Fusarium oxysporum* f. sp. *dianthi*, 742

d'Urso, F., Ayllón, M.A., Rubio, L., Sambade, A., Hermoso De Mendoza, A., Guerri, J. & Moreno, P. Contribution of uneven distribution of genomic RNA variants of citrus tristeza virus (CTV) within the plant to changes in the viral population following aphid transmission, 288
Daayf, F. *see* Mahuku, G.
Daayf, F. *see* Ongena, M.
Dabek, A.J. *see* Cronjé, P.
Dabek, A.J. *see* Cronjé, P.
Dahal, G., Ortiz, R., Tenkouano, A., Hughes, J.d'A., Thottappilly, G., Vuylsteke, D. & Lockhart, B.E.L. Relationship between natural occurrence of banana streak badnavirus and symptom expression, relative concentration of viral antigen, and yield

© 2000 BSPP

characteristics of some micropropagated *Musa* spp., 68
Danieli, B. *see* Curir, P.
Dann, E.K. & Deverall, B.J. Activation of systematic disease resistance in pea by an avirulent bacterium or a benzothiadiazole, but not by a fungal leaf spot pathogen, 324
Davies, D.L. & Adams, A.N. European stone fruit yellows phytoplasmas associated with a decline disease of apricot in southern England, 635
Davies, D.L. The occurence of two phytoplasmas associated with stunted *Rubus* species in the UK, 86
Dawson, W.A.J.M. & Bateman, G.L. Sensitivity of fungi from cereal roots to fluquinconazole and their suppressiveness towards take-all on plants with or without fluquinconazole seed treatment in a controlled environment, 477
De Haan, L.A.M., Numansen, A., Roebroeck, E.J.A. & Van Doorn, J. PCR detection of *Fusarium oxysporum* f.sp. *gladioli* race 1, causal agent of *Gladiolus* yellows disease, from infected corms, 89
De Lapeyre De Bellaire, L., Chillet, M., Dubois, C. & Mourichon, X. Importance of different sources of inoculum and dispersal methods of conida of *Colletotrichum musae*, the causal agent of banana anthracnose, for fruit contamination, 782
Dechamp-Guillaume, G. *see* Roustaee, A.
Dechamp-Guillaume, G. *see* Roustaee, A.
Delibes, A. *see* Andrés, M.F.
Desjardins, A.E. *see* Carter, J.P.
Deverall, B.J. *see* Colson-Hanks, E.S.
Deverall, B.J. *see* Dann, E.K.
Diao, A. *see* Chen, J.
Dickinson, M.J. *see* Mpunami, A.
Dodd, J.C. *see* Estrada, A.B.
Dolci, M. *see* Curir, P.
Du, J. *see* Chen, J.
Dubois, C. *see* De Lapeyre De Bellaire, L.
Dugdale, L.J., Mortimer, A.M., Isaac, S. & Collin, H.A. Disease response of carrot and carrot somaclones to *Alternaria dauci*, 57
Duggal, V., Jellis, G.J., Hollins, T.W. & Stratford, R. Resistance to powdery mildew in mutant lines of the susceptible wheat cultivar Hobbit 'sib', 468
Dumortier, V. *see* Lalaoui, F.
Duque, M.C. *see* Restrepo, S.
Duveiller, E. & García Altamirano, I. Pathogenicity of *Bipolaris sorokiniana* isolates from wheat roots, leaves and grains in Mexico, 235

Eden, M.A., Hill, R.A. & Galpoththage, M. An efficient baiting assay for quantification of *Phytophthora cinnamomi* in soil, 515
Egan, B. *see* Tran-Nguyen, L.
Elphinstone, J.G. *see* Pradhanang, P.M.
Elphinstone, J.G. *see* Pradhanang, P.M.
El-Tarabily, K.A., Soliman, M.H., Nassar, A.H., Al-Hassani, H.A., Sivasithamparam, K., McKenna, F. & Hardy, G.E.St.J. Biological control of *Sclerotinia minor* using a chitinolytic bacterium and actinomycetes, 573
Erselius, L.J. *see* Vega-Sánchez, M.E.
Estrada, A.B., Dodd, J.C. & Jeffries, P. Effect of humidity and temperature on conidial germination and appressorium development of two Philippine isolates of the mango anthracnose pathogen *Colletotrichum gloeosporioides*, 608
Evans, N. *see* West, J.S.
Evans, N. *see* Zhou, Y.

Fallik, E., Aharoni, Y., Copel, A., Rodov, V., Tuvia-Alkalai, S., Horev, B., Yekutieli, O., Wiseblum, A. & Regev, R. Reduction of postharvest losses of Galia melon by a short hot-water rinse, 333
Fernandez, D. *see* Plyler, T.R.
Fernandez, L. *see* Willocquet, L.
Fernández-Pavía, S. *see* Adorada, D.L.
Fitt, B.D.L. *see* Gilles, T.
Fitt, B.D.L. *see* Zhou, Y.
Flier, W.G. *see* Turkensteen, L.J.
Forbes, G.A. *see* Hijmans, R.J.
Forbes, G.A. *see* Vega-Sánchez, M.E.
Fox, R.T.V. *see* Pradhanang, P.M.
Fox, R.T.V. *see* Pradhanang, P.M.
Fox, R.T.V. *see* West, J.S.
Fry, W.E. *see* Vega-Sánchez, M.E.

Galal, A.A. *see* Ádám, A.L.
Galipienso, L., Navarro, L., Ballester-Olmos, J.F., Pina, J.A., Moreno, P. & Guerri, J. Host range and symptomatology of a graft-transmissible pathogen causing bud union crease of citrus on trifoliate rootstocks, 308
Galpoththage, M. *see* Eden, M.A.
García Altamirano, I. *see* Duveiller, E.
Gibb, K. *see* Padovan, A.
Gibb, K.S. *see* Tran-Nguyen, L.
Gilles, T., Fitt, B.D.L., Kennedy, R., Welham, S.J. & Jeger, M.J. Effects of temperature and wetness duration on conidial infection, latent period and asexual sporulation of *Pyrenopeziza brassicae* on leaves of oilseed rape, 498
Gilligan, C.A. *see* Webb, C.R.
Gladders, P. *see* Zhou, Y.
Goto, I. *see* Murakami, H.
Greif, C. *see* Seddas, A.
Guerri, J. *see* d'Urso, F.
Guerri, J. *see* Galipienso, L.
Guglieri, L. *see* Curir, P.
Guy, P.L. *see* Clarke, S.F.

Hafskjold Nærstad, R. *see* Hermansen, A.
Haidar, M.M. *see* Seddas, A.
Halama, P. *see* Lalaoui, F.
Halsey, K. *see* Mutasa-Gottgens, E.S.
Hannukkala, A. *see* Hermansen, A.
Hardy, G.E.St.J. *see* El-Tarabily, K.A.
Hardy, G.E.St.J. *see* Jackson, T.J.
Hargreaves, D. *see* Weller, S.A.
Harju, V.A. *see* Mumford, R.A.
Harvey, P.R., Butterworth, P.J., Hawke, B.G. & Pankhurst, C.E. Genetic variation among populations of *Pythium irregulare* in southern Australia, 619
Hashiba, T. *see* Narisawa, K.
Hawke, B.G. *see* Harvey, P.R.
Hélias, V., Andrivon, D. & Jouan, B. Development of symptoms caused by *Erwinia carotovora* ssp. *atroseptica* under field conditions and their effects on the yield of individual potato plants, 23
Hélias, V., Andrivon, D. & Jouan, B. Internal colonization pathways of potato plants by *Erwinia carotovora* ssp. *atroseptica*, 33
Henricot, B., Pérez Sierra, A. & Prior, C. A new blight disease on *Buxus* in the UK caused by the fungus *Cylindrocladium*, 805
Henry, C.M. *see* Mumford, R.A.
Hermansen, A., Hannukkala, A., Hafskjold Nærstad, R. & Brurberg, M.B. Variation in populations of *Phytophthora infestans* in Finland and Norway: mating type, metalaxyl resistance and virulence phenotype, 11
Hermoso De Mendoza, A. *see* d'Urso, F.
Hernández-Dorrego, A. *see* Andrés, M.F.
Hijmans, R.J., Forbes, G.A. & Walker, T.S. Estimating the global severity of potato late blight with GIS-linked disease forcast models, 697
Hill, R.A. *see* Eden, M.A.
Hille, J. *see* Mesbah, L.A.
Hoffer, M.E. *see* Cowger, C.
Hohl, H.R. *see* Vega-Sánchez, M.E.
Hollins, T.W. *see* Duggal, V.
Holt, J. *see* Zhang, X.-S.
Hooker, J.E. *see* Vigo, C.
Horev, B. *see* Fallik, E.
Howard, R.J. *see* Wang, H.
Hsiang, T., Ma, X.L., Yang, L. & Cook, S. Analyses of RAPD data for detection of host specialization in *Sclerotinia homoeocarpa*, 269
Hsu, Y.H., Annamalai, P., Lin, C.S., Chen, Y.Y., Chang, W.C. & Lin, N.S. A sensitive method for detecting bamboo mosaic virus (BaMV) and establishment of BaMV-free meristem-tip cultures, 101

Hu, B. *see* West, J.S.
Hughes, C. *see* West, J.S.
Hughes, J.d'A. *see* Dahal, G.
Hwang, S.F. *see* Wang, H.
Hyakumachi, M. *see* Pascual, C.B

Ioannidis, P.M. *see* Karaoglanidis, G.S.
Isaac, S. *see* Dugdale, L.J.

Jackson, T.J., Burgess, T., Colquhoun, I. & Hardy, G.E.StJ. Action of the fungicide phosphite on *Eucalyptus marginata* inoculated with *Phytophthora cinnamomi*, 147
Jacques, P. *see* Ongena, M.
Jacquet, C. *see* Seddas, A.
Jameson, P.E. *see* Clarke, S.F.
Jeffries, P. *see* Estrada, A.B.
Jeger, M.J. *see* Gilles, T.
Jeger, M.J. British Society for Plant Pathology Presidential Address 1999. Theory and Plant Epidemiology, 651
Jellis, G.J. *see* Duggal, V.
Johnson, R. Editorial, 1
Jones, P. *see* Cronjé, P.
Jones, P. *see* Cronjé, P.
Jones, P. *see* Mpunami, A.
Jouan, B. *see* Bouchek-Mechiche, K.
Jouan, B. *see* Hélias, V.
Jouan, B. *see* Hélias, V.
Julian, A.M. *see* Ngugi, H.K.
Jung, T., Blaschke, H. & Oßwald, W. Involvement of soilborne *Phytophthora* species in Central European oak decline and the effect of site factors on the disease, 706

Kalogerakis, S.N. *see* Vloutoglou, I.
Karaoglanidis, G.S., Ioannidis, P.M. & Thanassoulopoulos, C.C. Reduced sensitivity of *Cercospora beticola* isolates to sterol-demethylation-inhibiting fungicides, 567
Kennedy, R. *see* Clarkson, J.P.
Kennedy, R. *see* Gilles, T.
King, S.B. *see* Ngugi, H.K.
Kistler, H.C. *see* Plyler, T.R.
Kubelková, D. *see* Spak, J.

Lalaoui, F., Halama, P., Dumortier, V. & Paul, B. Cell wall-degrading enzymes produced *in vitro* by isolates of *Phaeosphaeria nodorum* differing in aggressiveness, 727
Lenzi, R. *see* Roggero, P.
Lesigne, P. *see* Roustaee, A.
Liddell, C.M. *see* Adorada, D.L.
Lin, C.S. *see* Hsu, Y.H.
Lin, N.S. *see* Hsu, Y.H.
Liu, S. *see* West, J.S.
Lobo Jnr, M., Lopes, C.A. & Silva, W.L.C. Sclerotinia rot losses in processing tomatoes grown under centre pivot irrigation in central Brazil, 51
Lockhart, B.E.L. *see* Dahal, G.
Lockley, D. *see* Paveley, N.D.
Lopes, C.A. *see* Lobo Jnr, M.
Lundsgaard, T. *see* Madriz-Ordeñana, K.

Ma, X.L. *see* Hsiang, T.
Madriz-Ordeñana, K., Rojas-Montero, R., Lundsgaard, T., Ramírez, P., Thordal-Christensen, H. & Collinge, D.B. Mechanical transmission of maize rayado fino marafivirus (MRFV) to maize and barley by means of the vascular puncture technique, 302
Mahuku, G., Peters, R.D., Platt, H.W. & Daayf, F. Random amplified polymorphic DNA (RAPD) analysis of *Phytophthora infestans* isolates collected in Canada during 1994 to 1996, 252
Manninger, K. *see* Ádám, A.L.
McIntosh, R.A. *see* Wellings, C.R.
McKenna, F. *see* El-Tarabily, K.A.
McPherson, G.M. *see* Weller, S.A.
Melgarejo, P. *see* Alfonso, C.
Mesbah, L.A., Van Der Weerden, G.M., Nijkamp, H.J.J. & Hille, J. Sensitivity among species of Solanaceae to AAL toxins produced by *Alternaria alternata* f. sp. *lycopersici*, 734
Metcalfe, R.J., Shaw, M.W. & Russell, P.E. The effect of dose and mobility on the strength of selection for DMI fungicide resistance in inoculated field experiments, 546
Moreno, M.T. *see* Sillero, J.C.
Moreno, P. *see* d'Urso, F.
Moreno, P. *see* Galipienso, L.
Moricca, S., Børja, I., Vendramin, G.G. & Raddi, P. Differentiation of *Seiridium* species associated with virulent cankers on cypress in the Mediterranean region by PCR-SSCP, 774
Morley, P.S. *see* Weller, S.A.
Mortimer, A.M. *see* Dugdale, L.J.
Mourichon, X. *see* De Lapeyre De Bellaire, L.
Mpunami, A., Tymon, A., Jones, P. & Dickinson, M.J. Identification of potential vectors of the coconut lethal disease phytoplasma, 355
Mukalazi, J. *see* Vega-Sánchez, M.E.
Mulder, A. *see* Turkensteen, L.J.
Mumford, R.A., Potyondi, L., Harju, V.A. & Henry, C.M. The identification of a phytoplasma from the Aster Yellows group infecting sugar beet in Hungary, 806
Mundt, C.C. *see* Cowger, C.
Murakami, H., Tsushima, S., Akimoto, T., Murakami, K., Goto, I. & Shishido, Y. Effects of growing leafy daikon (*Raphanus sativus*) on populations of *Plasmodiophora brassicae* (clubroot), 584
Murakami, K. *see* Murakami, H.
Murray, G.M. *see* Wellings, C.R.
Mutasa-Gottgens, E.S., Chwarszczynska, D.M., Halsey, K. & Asher, M.J.C. Specific polyclonal antibodies for the obligate plant parasite *Polymyxa* – a targeted recombinant DNA approach, 276

Narisawa, K., Ohki, K.T. & Hashiba, T. Suppression of clubroot and *Verticillium* yellows in Chinese cabbage in the field by the root endophytic fungus, *Heteroconium chaetospira*, 141
Nassar, A.H. *see* El-Tarabily, K.A.
Navarro, L. *see* Galipienso, L.
Newton, A.C. *see* Caten, C.E.
Ney, B. *see* Béasse, C.
Ngugi, H.K., Julian, A.M., King, S.B. & Peacocke, B.J. Epidemiology of sorghum anthracnose (*Colletotrichum sublineolum*) and leaf blight (*Exserohilum turcicum*) in Kenya, 129
Nicholson, P. *see* Carter, J.P.
Nicholson, P. *see* Simpson, D.R.
Nielsen, B.J. *see* O'Hara, R.B.
Nijkamp, H.J.J. *see* Mesbah, L.A.
Norman, J.R. *see* Vigo, C.
Numansen, A. *see* De Haan, L.A.M.
Nuñez, C. *see* Bertrand, B.

O'Hara, R.B., Nielsen, B.J. & Østergård, H. The effect of fungicide dose on the composition of laboratory populations of barley powdery mildew, 558
O'Neill, T.M. *see* Weller, S.A.
O'Neill, T.M. *see* Weller, S.A.
Ohki, K.T. *see* Narisawa, K.
Ojiambo, P.S. *see* Vega-Sánchez, M.E.
Ongena, M., Daayf, F., Jacques, P., Thonart, P., Benhamou, N., Paulitz, T.C. & Bélanger, R.R. Systematic induction of phytoalexins in cucumber in response to treatments with fluorescent pseudomonads, 523
Ortiz, R. *see* Dahal, G.
Oßwald, W. *see* Jung, T.
Østergård, H. *see* O'Hara, R.B.

Padovan, A., Gibb, K. & Persley, D. Association of '*Candidatus* Phytoplasma australiense' with green petal and lethal yellows diseases in strawberry, 362
Pankhurst, C.E. *see* Harvey, P.R.
Parry, D.W. *see* Simpson, D.R.
Pasco, C. *see* Bouchek-Mechiche, K.y
Pascual, C.B, Toda, T., Raymondo, A.D. & Hyakumachi, M. Characterization by conventional techniques and PCR of *Rhizoctonia solani* isolates causing banded leaf sheath blight

in maize, 108
Pasini, C. *see* Curir, P.
Paul, B. *see* Lalaoui, F.
Paulitz, T.C. *see* Ongena, M.
Paveley, N.D., Lockley, D., Vaughan, T.B., Thomas, J. & Schmidt, K. Predicting effective fungicide doses through observation of leaf emergence, 748
Peacocke, B.J. *see* Ngugi, H.K.
Peng, L. *see* West, J.S.
Pérez Sierra, A. *see* Henricot, B.
Persley, D. *see* Padovan, A.
Peters, R.D. *see* Mahuku, G.
Phelps, K. *see* Clarkson, J.P.
Pina, J.A. *see* Galipienso, L.
Pinochet, J. *see* Andrés, M.F.
Platt, H.W. *see* Mahuku, G.
Plyler, T.R., Simone, G.W., Fernandez, D. & Kistler, H.C. Genetic diversity among isolates of *Fusarium oxysporum* f.sp. *canariensis*, 155
Ponz, F. *see* Aguilar, I.
Potyondi, L. *see* Mumford, R.A.
Pradhanang, P.M., Elphinstone, J.G. & Fox, R.T.V. Identification of crop and weed hosts of *Ralstonia solanacearum* biovar 2 in the hills of Nepal, 403
Pradhanang, P.M., Elphinstone, J.G. & Fox, R.T.V. Sensitive detection of *Ralstonia solanacearum* in soil: a comparison of different detection techniques, 414
Price, T.V. *see* Suheri, H.
Prior, C. *see* Henricot, B.

Raddi, P. *see* Moricca, S.
Raggi, V. Hydroxyproline-rich glycoprotein accumulation in tobacco leaves protected against *Erysiphe cichoracearum* by potato virus Y infection, 179
Raman, H. *see* Wellings, C.R.
Ramírez, P. *see* Madriz-Ordeñana, K.
Raposo, R. *see* Alfonso, C.
Raymondo, A.D. *see* Pascual, C.B
Regev, R. *see* Fallik, E.
Restrepo, S., Duque, M.C. & Verdier, V. Charcteriazation of pathotypes among isolates of *Xanthomonas axonopodis* pv. *manihotis* in Colombia, 680
Rezanoor, H.N. *see* Carter, J.P.
Rezanoor, H.N. *see* Simpson, D.R.
Ridout, M.S. *see* Xu, X.-M.
Robinson, J.D. *see* Xu, X.-M.
Robinson, J.D. *see* Xu, X-M.
Rodgers-Gray, B.S. & Shaw, M.W. Substantial reductions in winter wheat diseases caused by addition of straw but not manure to soil, 590
Rodov, V. *see* Fallik, E.
Rodriguez, A.M. *see* Vega-Sánchez, M.E.
Roebroeck, E.J.A. *see* De Haan, L.A.M.
Roggero, P., Ciuffo, M. & Lenzi, R. Natural infection by *Cucumber mosaic virus* (*Cucumovirus*; *Bromoviridae*) in *Fuchsia boliviana Carr* and the insuseptibility of other *Fuchsia* species and hybrids to the virus, 802
Rojas-Montero, R. *see* Madriz-Ordeñana, K.
Roustaee, A., Barrault, G., Dechamp-Guillaume, G., Lesigne, P. & Sarrafi, A. Inheritance of partial resistance to black stem (*Phoma macdonaldii*) in sunflower, 396
Roustaee, A., Costes, S., Dechamp-Guillaume, G. & Barrault, G. Phenotypic variabilty of *Leptosphaeria lindquistii* (anamorph: *Phoma macdonaldii*), a fungal pathogen of sunflower, 227
Rovira, M. *see* Aramburu, J.M.
Rubiales, D. *see* Sillero, J.C.
Rubio, L. *see* d'Urso, F.
Russell, P.E. *see* Metcalfe, R.J.

Sacco, M. *see* Curir, P.
Sambade, A. *see* d'Urso, F.
Sánchez, F. *see* Aguilar, I.
Santini, A. & Capretti, P. Analysis of the Italian population of *Ceratocystis fimbriata* f.sp. *platani* using RAPD and minisatellite markers, 461
Sarah, J.-L. *see* Bertrand, B.
Sarrafi, A. *see* Roustaee, A.
Savary, S. *see* Willocquet, L.
Schmidt, K. *see* Paveley, N.D.
Seddas, A., Haidar, M.M., Greif, C., Jacquet, C., Cloquemin, G. & Walter, B. Establishment of a relationship between grapevine leafroll closteroviruses 1 and 3 by use of monoclonal antibodies, 80
Shaw, M.W. *see* Metcalfe, R.J.
Shaw, M.W. *see* Rodgers-Gray, B.S.
Shishido, Y. *see* Murakami, H.
Sillero, J.C., Moreno, M.T. & Rubiales, D. Characterization of new sources of resistance to *Uromyces viciae-fabae* in a germplasm collection of *Vicia faba*, 389
Silva, W.L.C. *see* Lobo Jnr, M.
Simone, G.W. *see* Plyler, T.R.
Simpson, D.R., Rezanoor, H.N., Parry, D.W. & Nicholson, P. Evidence for differential host preference in *Microdochium nivale* var. *majus* and *Microdochium nivale* var. *nivale*, 261
Sivasithamparam, K. *see* El-Tarabily, K.A.
Soliman, M.H. *see* El-Tarabily, K.A.
Spak, J. & Kubelková, D. Serological variability among European isolates of radish mosaic virus, 295
Spence, N.J. *see* Brunt, A.A.
Stead, D.E. *see* Weller, S.A.
Stead, D.E. *see* Weller, S.A.
Stratford, R. *see* Duggal, V.
Suheri, H. & Price, T.V. Infection of onion leaves by *Alternaria porri* and *Stemphylium vesicarium* and disease development in controlled environments, 375

Tenkouano, A. *see* Dahal, G.
Thanassoulopoulos, C.C. *see* Karaoglanidis, G.S.
Thomas, J. *see* Paveley, N.D.
Thonart, P. *see* Ongena, M.
Thordal-Christensen, H. *see* Madriz-Ordeñana, K.
Thottappilly, G. *see* Dahal, G.
Thrall, P.H. & Burdon, J.J. Effect of resistance variation in a natural plant host–pathogen metapopulation on disease dynamics, 767
Tivoli, B. *see* Béasse, C.
Toda, T. *see* Pascual, C.B
Tran-Nguyen, L., Blanche, K.R., Egan, B. & Gibb, K.S. Diversity of phytoplasmas in northern Australian sugarcane and other grasses, 666
Tsushima, S. *see* Murakami, H.
Turkensteen, L.J., Flier, W.G., Wanningen, R. & Mulder, A. Production, survival and infectivity of oospores of *Phytophthora infestans*, 688
Turnbull, G.D. *see* Wang, H.
Tuvia-Alkalai, S. *see* Fallik, E.
Tymon, A. *see* Mpunami, A.
Tymon, A.M. *see* Cronjé, P.
Tymon, A.M. *see* Cronjé, P.

Van Der Weerden, G.M. *see* Mesbah, L.A.
Van Doorn, J. *see* De Haan, L.A.M.
Vaughan, T.B. *see* Paveley, N.D.
Vega-Sánchez, M.E., Erselius, L.J., Rodriguez, A.M., Bastidas, O., Hohl, H.R., Ojiambo, P.S., Mukalazi, J., Vermeulen, T., Fry, W.E. & Forbes, G.A. Host adaptation to potato and tomato within the US–1 clonal lineage of *Phytophthora infestans* in Uganda and Kenya, 531
Vendramin, G.G. *see* Moricca, S.
Verdier, V. *see* Restrepo, S.
Vermeulen, T. *see* Vega-Sánchez, M.E.
Vigo, C., Norman, J.R. & Hooker, J.E. Biocontrol of the pathogen *Phytophthora parasitica* by arbuscular mycorrhizal fungi is a consequence of effects on infection loci, 509
Vloutoglou, I. & Kalogerakis, S.N. Effects of inoculum concentration, wetness duration and plant age on development of early blight (*Alternaria solani*) and on shedding of leaves in tomato plants, 339
Vuylsteke, D. *see* Dahal, G.

Walker, T.S. *see* Hijmans, R.J.

Wallwork, H. *see* Wellings, C.R.
Walter, B. *see* Seddas, A.
Wang, H., Hwang, S.F., Chang, K.F., Turnbull, G.D. & Howard, R.J. Characterization of *Ascochyta* isolates and susceptibility of pea cultivars to the ascochyta disease complex in Alberta, 540
Wanningen, R. *see* Turkensteen, L.J.
Waugh, K.O. *see* Adorada, D.L.
Waugh, M.E. *see* Adorada, D.L.
Webb, C.R., Gilligan, C.A. & Asher, M.J.C. Modelling the effect of temperature on the development of *Polymyxa betae*, 600
Welham, S.J. *see* Gilles, T.
Welham, S.J. *see* Zhou, Y.
Weller, S.A., Stead, D.E., O'Neill, T.M. & Morley, P.S. Root mat of tomato caused by rhizogenic strains of *Agrobacterium* biovar 1 in the UK, 799
Weller, S.A., Stead, D.E., O'Neill, T.M., Hargreaves, D. & McPherson, G.M. Rhizogenic *Agrobacterium* biovar 1 and cucumber root mat in the UK, 43
Wellings, C.R., Burdon, J.J., McIntosh, R.A., Wallwork, H., Raman, H. & Murray, G.M. A new variant of *Puccinia striiformis* causing stripe rust on barley and wild *Hordeum* species in Australia, 803
West, J.S., Evans, N., Liu, S., Hu, B. & Peng, L. *Leptosphaeria maculans* causing stem canker of oilseed rape in China, 800
West, J.S., Hughes, C. & Fox, R.T.V. *Armillaria mellea* can infect the perennial weed, *Rumex obtusifolius*, in the UK, 808
Willocquet, L., Fernandez, L. & Savary, S. Effect of various crop establishment methods practised by Asian farmers on epidemics of rice sheath blight caused by *Rhizoctonia solani*, 346
Wiseblum, A. *see* Fallik, E.

Xu, X.-M. & Ridout, M.S. Stochastic simulation of the spread of race-specific and race-nonspecific aerial fungal pathogens in cultivar mixtures, 207
Xu, X.-M. & Robinson, J.D. Epidemiology of brown rot (*Monilinia fructigena*) on apple: infection of fruits by conidia, 201
Xu, X-M. & Robinson, J.D. Effects of temperature on the incubation and latent periods of hawthorn powdery mildew *(Podosphaera clandestina)*, 791

Yang, J.-P. *see* Chen, J.
Yang, L. *see* Hsiang, T.
Yekutieli, O. *see* Fallik, E.

Zhang, X.-S., Holt, J. & Colvin, J. A general model of plant-virus disease infection incorporating vector aggregation, 435
Zhou, Y., Fitt, B.D.L., Welham, S.J., Evans, N. & Gladders, P. Effects of stem canker (*Leptosphaeria maculans*) and light leaf spot (*Pyrenopeziza brassicae*) on yield of winter oilseed rape (*Brassica napus*) in southern England, 487

Keyword index

AAL toxins, 734
ACC, 428
acetophenone, 742
active oxygen species, 317
adaptation, 531
aetiology, 23
aggressiveness, 227, 727
Agrobacterium, 43
agroecological zones, 697
air drying, 515
Allium, 375
allopurinol, 317
allozymes, 252
Alternaria dauci, 57
Alternaria macrospora, 171
Alternaria porri, 375
Alternaria solani, 339
Alternaria, 734
amino acid sequence, 370
analysing epidemics, 748
anastomosis groups, 108
anthracnose, 129, 608
ApMv transmission, 423
apple, 201
appressoria, 375, 608
apricot, 635
Arabidopsis, 659
Ascochyta, 540
ascospores, 227
AUDPC, 207

baiting assay systems, 515
bamboo mosaic potexvirus, 101
banana streak badnavirus, 68
banana, 68
barley powdery mildew, 558
barley, 302
Bemisia tabaci, 435
benzothiadiazole, 171, 324
bioassay, 89
biocontrol, 141
biological control, 509
biological indexing, 308
biovar 2, 43, 403
bitertanol, 567
black stem, 396
blackleg, 23
blackstem, 227
BNYVV, 600
Botrytis squamosa, 119
brassicas, 295
brushes, 333
BSV, 68

Camelina sativa, 295
canker strain, 461
Capsicum annuum, 719
carnation, 742
carrot, 57
cassava bacterial blight, 680
cassava mosaic virus disease, 435
cell-wall-degrading enzymes, 573
Ceratocystis fimbriata, 461
Cercospora leaf spot, 567
cereals, 261, 477
chlorophyll, 57
citrange rootstock, 308
citrus virus, 308
closterovirus, 80
clubroot, 141
cluster analysis, 108
Cochliobolus sativus, 235
coconut, 355
Coffea arabica, 383
Colletotrichum, 608
Colletotrichum, 782
colony size, 389
combined yield loss, 487
combining ability, 396
common scab, 3
comovirus, 295
competition, 261
complementary DNA (cDNA), 288
conidial production, 498
contact rate, 435
contamination, 782
control of wheat leaf rust, 317
corky root disease, 383
Corylus avellana, 423
cotton, 171
crazy top, 628
cross-protection, 659
cross-reactivity, 80
CTV isolates, 288
cucumber root mat, 43
Cucumis sativus, 523
cultivar, 498
cultivar resistance, 33
cultivar response, 370
cypress canker, 773

Dalbulus maidis, 302
DAS-ELISA, 33
Daucus carota, 57
decoy plant, 584
defoliation, 339
detection, 414
detoxification, 742
developing countries, 697
diagnostic markers, 640
Diastrombus mkurangai, 355
2,6-dichloroisonicotinic acid, 171
Dig-labelled probe, 101
dihydrozeatin, 428
direct seeding, 346
disease bioassay, 57
disease complex, 383
disease forecasting, 498
disease intensity, 51
disease progress curves, 129, 498
disease severity, 584
disease spread, 346
dispersal, 782
DNA fingerprinting, 155
double-stranded RNA (dsRNA), 288
dsRNA, 428

ecology, 3
ELISA, 414, 423
elm, 86
England, 635
environmental variation, 558
enzymes, 727
epidemic development, 51
epidemiology, 33, 43, 129, 295, 435, 600, 608, 688, 782
Erysiphe cichoracearum, 179
Erysiphe graminis, 468, 590
evolution, 619
excised leaf assay, 540

faba-bean, 389
fenpropimorph, 558, 748

field populations, 219
fluorescence microscopy, 584
fluorescent pseudomonads, 523
fluquinconazole, 477, 546
flutriafol, 546, 567
foliar blight, 235
fruit maturity, 201
fruit, 635
fungal antagonism, 477
fungal population, 461
fungicide resistance, 558, 567
fungicides, 697
Fusarium oxysporum f.sp. *dianthi*, 742
Fusarium oxysporum f.sp. *gladioli*, 89
Fusarium oxysporum, 383
Fusarium, 155, 590

Gaeumannomyces graminis, 477
gene-for-gene, 767
genetic diversity, 155
genetic variation, 243
germination, 119, 608
Gibberella zeae, 452
Gladiolus, 89
glume blotch of wheat, 219
glutathione-s-transferase, 276
graft inoculation, 308
grapevine, 80
grassy shoot, 666
grey mould, 243

hawthorn, 791
head blight, 452
Helianthus annuus, 227
Heteroconium chaetospira, 141
heterogeneity, 362
His-Tag fusion protein, 276
host defence, 147
host range, 403
host resistance, 697
host–pathogen interaction, 235
hot water rinse, 333
hybridoma, 80
hydroponic crops, 43
hydroxamic acid, 261
hydroxyproline-rich glycoprotein, 179
hypersensitive response, 317
hypersensitivity, 389
hyphal development, 468
hyphal lysis, 573

immunoblotting, 80
immunogold labelling, 80
inculation period, 791
indicator plants, 308
induced resistance, 590
induced systemic resistance, 523
infection criteria, 498
infection studies, 719
infection, 201
inoculum concentration, 339
inoculum sources, 781
intraspecific diversity, 640
isoeclectrofocusing, 640
isozyme markers, 640
ITS2 sequence variation, 774

jarrah, 147
jasmonic acid, 428

late blight, 688
latent infection, 33
latent period, 389, 791
LD phytoplasmas, 355
leaf layers, 748
leaf wetness, 339, 375
leafroll, 80
leaf-spot disease, 57
Leptosphaeria maculans, 487
lettuce microbial antagonists, 573
light leaf spot, 487, 498
lineages, 531
Linum marginale, 767
Lycopersicon esculentum, 509
Lycopersicon, 734

maize, 452
maize downy mildew, 628
maize rayado fino virus, 302
majus, 261
mango, 608
marketing, 333
mathematical model, 435
mating type, 11, 252
mechanical transmission, 302
Meenoplus spp., 355
Melampsora lini, 767
Meloidogyne arabicida, 383
meristem culture, 101
metalaxyl resistance, 11
metalaxyl sensitivity, 252
Microdochium nivale, 261
minisatellite markers, 461
mixture, 207
modelling, 600, 791
molecular markers, 269
Monilinia fructigena, 201
multiplex PCR, 89
Musa, 68, 782
mustard, 295
mutant lines resistance, 468
mycorrhiza, 509
Mycosphaerella graminicola, 445, 546, 590
Mycosphaerella pinodes, 324

necrosis, 389, 734
Nepal, 403
netted scab, 3
nucleotide sequence, 370

oospores, 628
ORMV, 659
outcrossing, 619
overwintering, 403

P. betae, 600
palm wilt, 155
parameter estimates, 129
partial resistance, 396
pathogenic variability, 235
pathogenicity, 3
pathotypes, 680
PCR, 33, 243, 414, 540
PCR products, 355
PCR-RFLP analysis, 108
pea, 324, 540
peach, 86
Peronosclerospora sorghi, 628
peroxidase, 719
Phaeosphaeria nodorum, 727
phenolics, 147
Phoenix canariensis, 155
Phoma macdonaldii, 396
phosphonate, 147
photosynthesis, 187
physiologic specialization, 445
physiological specialization, 269
phytoalexins, 523
phytoanticipin, 742
Phytophthora cinnamomi, 515
Phytophthora infestans, 11, 252, 531, 697
Phytophthora, 147, 706
phytoplasma, 362, 666
phytoplasmas, 86, 635
plant age, 339
plant growth-promoting rhizobacteria, 523
plantain, 68
planting date, 129
Plasmodiophora brassicae, 276
Platonus acerifolia, 461

plochloraz, 546
polymerase chain reaction-single strand conformation polymorphism (PCR-SSCP), 773
Polymyxa graminis, 276
population genetics, 619
population structure, 243
populations, 531
postharvest disease, 608
potato late blight, 11
potato late blight, 252
potato virus Y, 179
potato, 3, 33, 531, 688
powdery mildew, 791
PR proteins, 324
Pratylenchus, 640
premoistening, 515
progeny tubers, 33
propiconazole, 748
Pseudomonas syringae, 324
Puccinia recondita, 590
Puccinia striiformis, 748
pycniospores, 227
Pyrenopeziza brassicae, 487
Pythium root rot, 619

quantitative PCR, 261
Quercus, 706

race-nonspecific pathogen, 207
race-nonspecific resistance, 688
races, 89
race-specific pathogen, 207
radiation interception efficiency, 187
radiation use efficiency, 187
radioactive and nonradioactive hybridization, 101
Radopholus similus, 640
Ralstonia solanacearum, 403, 414
RAPD, 461, 540
RAPD polymorphism, 243
RAPDs, 89, 252
rate, 207
reduced doses, 546
resistance, 3, 540, 767
Rhizoctonia solani, 108
Rhizoctonia, 346
rhizomania, 276, 600
rhizosphere competence, 573
rhizosphere, 706
ribiprobe, 101
rice cultivation, 346
rice, 452
Ri-plasmid, 43
root architecture, 509
root colonization, 573
root endophytic fungus 141
root-lesion nematode, 640
RT-PCR, 288
Rubus, 86

salicic acid, 171
salicylic acid, 428
satellite RNA, 101
sclerotia, 119
Sclerotinia sclerotiorum, 51
screening for rust resistance, 389
seed treatment fungicide, 477
Seiridium cardinale, 774
Seiridium cupressi, 774
selection, 558
selective medium, 414
sensitivity, 734
Septoria nodorum, 219
septoria tritici blotch, 445
Septoria tritici, 546, 748
sexual reproduction, 219
sheath blight, 346
silicon nutrition, 590
simulation, 207
single-strand conformation polymorphism analysis (SSCP analysis), 288
soft rot, 23
soil, 414
soil particle size, 515
soil pH, 706
soil texture, 706
Solanum tuberosum, 23
somaclones, 57
sorghum cultivars, 129
sources of inoculum, 688
spatial structure, 767
species relatedness, 640
spore germination, 584
sporulation, 509
spot blotch, 235
16SrRNA, 362
Stagonospora nodorum, 727
stem canker, 487
Stemphylium vesicarium, 375
stems, 33
stolons, 33
stone, 635
storage, 333
strawberry, 362
Streptomyces, 3
sugar beet, 276, 567
sugarcane, 666
sunflower, 396
symptom expression, 68
systemic resistance, 171, 324
systemicity, 546

take-all, 477
taxonomy, 640
temperature, 119, 515
temperature effects, 600
thaumatin-like gene, 468
TMGMV, 659
tobacco, 179, 659
tobamovirus, 659
tomato, 339, 531, 688, 734
tomatoes, 51
transplanting, 346
Triticum aestivum, 445, 468
turfgrass disease, 269

vascular pressure, 302
vectors, 355
vegetative compatibility, 155
vegetative incompatibility, 219
Verticillium yellows, 141
virulence phenotype, 11
virulence, 540, 680, 767
virus transmission, 435

water potential, 119
weeds, 403
wheat, 235, 452
wheat yellow mosaic virus isolate, 370
white clover mosaic potexvirus, 428
whiteleaf, 666
winter bread wheat, 445
winter oilseed rape, 487
wound age, 201

X group, 86
Xanthomonas axonopodis pv. *manihotis*, 680
yellow leaf syndrome, 666
yellows, 635
yellows group, 86
yield loss, 23, 51

Zea mays, 108
zoospores, 719